

A global movement toward an ecosystem approach to management of marine resources

K. Sherman^{1,*}, M. Sissenwine², V. Christensen³, A. Duda⁴, G. Hempel⁵, C. Ibe⁶, S. Levin⁷, D. Lluch-Belda⁸, G. Matishov⁹, J. McGlade¹⁰, M. O'Toole¹¹, S. Seitzinger¹², R. Serra¹³, H.-R. Skjoldal¹⁴, Q. Tang¹⁵, J. Thulin¹⁶, V. Vandeweerdt¹⁷, K. Zwanenburg¹⁸

¹Northeast Fisheries Science Center, Narragansett Laboratory, NOAA-Fisheries, 28 Tarzwell Drive, Narragansett, Rhode Island 02882, USA

* Email: kenneth.sherman@noaa.gov

²Director of Scientific Programs, NOAA-Fisheries, Silver Spring, Maryland 20910-3282, USA

³Fisheries Centre, University of British Columbia, Vancouver, British Columbia, V6T 1Z4, Canada

⁴International Waters, Global Environment Facility (GEF) Secretariat, World Bank, Washington, DC 20433, USA

⁵Berater des Präsidenten des Senats für den Wissenschaftsstandort Bremen-Bremerhaven, Senate of Bremen, Tiefer 2, 28195 Bremen, Germany

⁶United Nations Industrial Development Organization (UNIDO), United Nations Compound, Accra, PO Box 1423, Ghana

⁷Department of Ecology and Evolutionary Biology, Princeton University, Princeton, New Jersey 08544-1003, USA

⁸Centro Interdisciplinario de Ciencias Marinas (CICIMAR), Instituto Politécnico Nacional (IPN), La Paz, Baja California Sur, 23096, Mexico

⁹Academy of Sciences, Murmansk Marine Biological Institute (MMBI), Azov Branch, Rostov-on-Don 344066, Russia

¹⁰European Environment Agency, Copenhagen K1050, Denmark

¹¹Benguela Current LME Programme, Coordination Unit, Windhoek, PO Box 40728, Namibia

¹²Institute of Marine & Coastal Sciences, Rutgers, State University of New Jersey, Cook Campus, New Brunswick, New Jersey 08901-8521, USA

¹³Instituto de Fomento Pesquero, IFOP, Valparaiso, Casilla 8-V, Chile

¹⁴Institute of Marine Research, Bergen 5024, PO Box 1870, Norway

¹⁵Yellow Sea Fisheries Research Institute, Qingdao 266071, China

¹⁶International Council for the Exploration of the Sea (ICES), Copenhagen K1261, Denmark

¹⁷United Nations Environment Programme (UNEP), PO Box 16277, 2500 BE, The Hague, Netherlands

¹⁸Marine Fish Division, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, B2Y 4A4, Canada

The Large Marine Ecosystem Approach. Reports of problems with marine ecosystems are widespread in the scientific literature and the news media. Calls for an ecosystem approach to resource assessment and management are seldom accompanied by a practical strategy, particularly one with a payment plan for the approach in developing countries. However, a global movement that makes the ecosystem approach to management practical already exists. It is known as the Large Marine Ecosystem (LME) approach, and it is being endorsed and supported by governments world-

wide, as well as by a broad constituency in the scientific community.

While we concur with the movement toward an ecosystem-based approach to the management of marine fisheries (Gislason & Sinclair 2000, Pitcher 2001, Stergiou 2002, Garcia et al. 2003, Sainsbury & Sumaila 2003, Browman et al. 2004, Pikitch et al. 2004), it is important to recognize that a broader, place-based approach to marine ecosystem assessment and management, focused on clearly delineated ecosystem units, is needed and is presently under way, with the support of financial grants, donor and UN partnerships, in nations of Africa, Asia, Latin America and eastern Europe. It is within the boundaries of 64 LMEs that (1) 90% of the world's annual yield of marine fisheries is produced (Garibaldi & Limongelli 2003), (2) global levels of primary production are the highest, (3) the degradation of marine habitats is most severe, and (4) coastal pollution is concentrated and levels of eutrophication are increasing (GESAMP 2001). Large marine ecosystems (LMEs) are natural regions of coastal ocean space encompassing waters from river basins and estuaries to the seaward boundaries of continental shelves and outer margins of coastal currents and water masses (cf. Fig. 4). They are relatively large regions characterized by distinct bathymetry, hydrography, productivity, and trophically dependent populations (Alexander 1990, Levin 1990, Sherman 1994; see www.edc.uri.edu/lme).

Since 1995, the Global Environment Facility (GEF) has provided substantial funding to support country-driven projects for introducing multisectoral ecosystem-based assessment and management practices for LMEs located around the margins of the oceans. At present, 121 developing countries are engaged in the preparation and implementation of GEF-LME projects, totaling \$650 million in start-up funding. A total of 10 projects including 70 countries has been approved by the GEF Council, and another 7 projects involving 51 countries have GEF international waters projects under preparation (see www.iwlearn.net).

A 5 module indicator approach to assessment and management of LMEs has proven useful in ecosystem-based projects in the USA and elsewhere, using suites of indicators of LME productivity, fish and fisheries, pollution and ecosystem health, socioeconomic, and governance. The productivity indicators include spatial and temporal measurements of temperature, salinity, oxygen, nutrients, primary productivity, chlorophyll, zooplankton biomass, and biodiversity. For fish and fisheries, indicators are catch and effort statistics, demersal and pelagic fish surveys, fish population demography, and stock assessments (NMFS 1999). Pollution and ecosystem health indicators include quality indices for water, sediment, benthos, habitats,




Fig. 4. Boundaries of the 64 Large Marine Ecosystems (LMEs) of the world and primary productivity ($\text{gC m}^{-2} \text{yr}^{-1}$). Annual productivity estimates are based on SeaWiFS satellite data collected between September 1998 and August 1999, and on the model developed by Behrenfeld & Falkowski (1997). Color-enhanced image provided by Rutgers University (available at: www.edc.uri.edu/lme, Introduction)

and fish tissue contaminants (EPA 2004). Socioeconomic and governance indicators are discussed in Sutinen et al. (2000) and Juda & Hennessey (2001). The modules are adapted to LME conditions through a transboundary diagnostic analysis (TDA) process, to identify key issues, and a strategic action program (SAP) development process for the groups of nations or states sharing an LME, to remediate the issues (Wang 2004). These processes are critical for integrating science into management in a practical way, and for establishing appropriate governance regimes. Of the 5 modules, 3 modules apply science-based indicators that focus on productivity, fish/fisheries, and pollution/ecosystem health, and the other 2 modules, socioeconomics and governance, focus on economic benefits to be gained from a more sustainable resource base and from providing stakeholders and stewardship interests with legal and administrative support for ecosystem-based management practices. The first 4 modules support the TDA process, while the governance module is associated with periodic updating of the SAP development process. Adaptive management regimes are encouraged through periodic assessment processes (TDA updates) and through updating the action programs as gaps are filled.

The GEF-LME projects presently funded or in the pipeline for funding in Africa, Asia, Latin America and eastern Europe represent a growing network of marine scientists, marine managers, and ministerial leaders

who are pursuing ecosystem and fishery recovery goals. The annual fisheries biomass yields from the ecosystems in the network are 44.8% of the global total, and are a firm basis for movement by the participating countries toward the 2002 World Summit on Sustainable Development (WSSD) targets for introducing ecosystem-based assessment and management by 2010, and for recovering depleted stocks and achieving fishing at maximum sustainable yield levels by 2015. The FAO Code of Conduct for Responsible Fisheries (FAO 1995) is supported by most coastal nations and has immediate applicability to reaching the WSSD fishery goals. The code argues for moving forward with a precautionary approach to fisheries sustainability, using available information more conservatively to err on the side of lower total allowable catch levels than has been the general practice in past decades. Although fishing effort data are not available in FAO global catch reporting statistics and could bias catch data interpretations, it appears that the biomass and yields of 11 species groups in 6 LMEs have been relatively stable or have shown marginal increases over the period from 1990 to 1999. The yield for these 6 LMEs—the Arabian Sea, Bay of Bengal, Indonesian Sea, North Brazil Shelf, Mediterranean Sea and the Sulu-Celebes Sea—was 8.1 million t, or 9.5% of the global marine fisheries yield in 1999 (Garibaldi & Limongelli 2003). The countries bordering these 6 LMEs are among the world's most populous, representing approximately one-quarter of

the total human population. These LME border countries increasingly depend on marine fisheries for food security, and for national and international trade. Given the risks of fishing down the food web, it would appear opportune for the stewardship agencies responsible for the fisheries of the LME-bordering countries to limit increases in fishing effort during a period of relative biomass stability.

Evidence for species biomass recovery following significant reduction in fishing effort through mandated actions is encouraging. In the USA Northeast Shelf LME, management actions to reduce fishing effort contributed to a recovery of depleted herring and mackerel stocks and an initiation of the recovery of depleted yellowtail flounder and haddock stocks (Sherman et al. 2003); this was in combination with the robust condition of average annual primary productivity ($350 \text{ g C m}^{-2} \text{ yr}^{-1}$) for the past 3 decades, a relatively stable zooplankton biomass at or near 33 cm^3 per 100 m^3 for the past 30 yr (Sherman et al. 2002), and an oceanographic regime marked by a recurring pattern of inter-annual variability, but showing no evidence of temperature shift of the magnitude described for other North Atlantic LMEs, including the Scotian Shelf (Zwanenburg 2003), the Newfoundland-Labrador Shelf (Rice 2002), the Iceland Shelf (Astthorsson & Vilhjálmsson 2002) and the North Sea (Perry et al. 2005). On the other hand, 3 LMEs remain at high risk for fisheries biomass recovery—expressed as a pre-1960s ratio of demersal to pelagic species—the Gulf of Thailand, East China Sea, and Yellow Sea (Pauly & Chuenpagdee 2003, Chen & Shen 1999, Tang & Jin 1999). The People's Republic of China has initiated steps toward recovery by mandating 60–90 d closures to fishing in the Yellow Sea and East China Sea (Tang 2003). The country-driven planning and implementation documents supporting the ecosystem approach to LME assessment and management practices can be found at www.iwlearn.net.

Nitrogen loadings. Globally, LME projects, in addition to rebuilding depleted fish stocks and restoring degraded coastal habitats, are also concerned with the mitigation of the effects of nitrogen loadings. Nitrogen over-enrichment has been a coastal problem for 2 decades in the Baltic Sea LME (HELCOM 2001). More recent human-induced increases in nitrogen flux range from 4- to 8-fold in the USA from the Gulf of Mexico to the New England coast (Howarth et al. 2000). In European LMEs, recent nitrogen flux increases have ranged from 3-fold in Spain to 11-fold in the Rhine River basin draining to the North Sea LME (Howarth et al. 2000). This disruption of the nitrogen cycle originated in the Green Revolution of the 1970s as the world community converted wetlands to agriculture, utilized more chemical fertilizer, and expanded

irrigation to feed the world (Duda & El-Ashry 2000). For the estuaries of the southeastern USA (Duda 1982) and for the Gulf of Mexico (Rabalais et al. 1999), much of the increase in nitrogen export to LMEs is from agricultural inputs, from the increased delivery of nitrogen fertilizer as wetlands were converted to agriculture, and from livestock production (NRC 2000). Also, sewage from large cities is a significant contributor to eutrophication, as is increased nitrogen in atmospheric deposition resulting from combustion of fossil fuels by automobiles and industrial activities (GESAMP 2001).

Global forecast models of nitrogen export from freshwater basins to coastal waters indicate that there will be a 50% increase world-wide in dissolved inorganic nitrogen (DIN) export by rivers to coastal systems from 1990 to 2050 (Seitzinger & Kroeze 1998, Kroeze & Seitzinger 1998). Such increases in nitrogen export are alarming for the future sustainability of LMEs. Given the expected future increases in population and in fertilizer use, without significant mitigation of nitrogen inputs, LMEs will be subjected to a future of increasing harmful algal bloom events, reduced fisheries, and hypoxia that further degrades marine biomass yields and biological diversity. Models of nitrogen loading from land-based sources and models of ecosystem structure and function are being applied to LMEs with financial assistance from the GEF. Estimates of carrying capacity using ECOPATH-ECOSIM food web approaches for the world's 64 LMEs are being prepared in a GEF-supported collaboration between scientists of the University of British Columbia and marine specialists from developing countries. Similarly, a 24 mo training project is being implemented by scientists from Rutgers University in collaboration with IOC/UNESCO to estimate expected nitrogen loadings for each LME over the next decade. Scientists from Princeton University and the University of California at Berkeley are examining particle spectra and pattern formation within LMEs. Additionally, the American Fisheries Society and the World Council of Fisheries Societies are collaborating in an electronic network to expedite information access and communication among marine specialists (for details on the GEF-LME project, see www.gefonline.org/projectDetails.cfm?projID=2474).

The growing number of country-driven commitments to move toward ecosystem-based assessment and management of marine resources and environments provides an unprecedented opportunity for accelerating the transition to sustainable use, conservation, and development of marine ecosystems. The social, economic, and environmental costs of inaction are simply too high for multilateral and bilateral institutions and international agencies not to support the initial efforts of 121 countries attempting to reach the WSSD marine ecosystem targets for restoration and sustainability. Both developed and developing nations

have a stake in moving toward the use of sustainable ecosystem resources. Momentum should not be lost, as this could result in irreversible damage to coastal ecosystems, to the livelihoods and security of poor coastal communities, and to the economies of coastal nations.

LITERATURE CITED

- Alexander LM (1990) Geographic perspectives in the management of large marine ecosystems. In: Sherman K, Alexander LM, Gold BD (eds) Large marine ecosystems: patterns, processes and yields. American Association for the Advancement of Science, Washington, DC, p 220–223
- Asthorsson OS, Vilhjálmsson H (2002) Iceland Shelf LME: decadal assessment and resource sustainability. In: Sherman K, Skjoldal HR (eds) Large Marine Ecosystems of the North Atlantic. Elsevier, Amsterdam, p 219–244
- Behrenfeld M, Falkowski PG (1997) Photosynthetic rates derived from satellite-based chlorophyll concentrations. *Limnol Oceanogr* 42(1):1–20
- Browman HI, Stergiou KI (eds) (2004) Perspectives on ecosystem-based approaches to the management of marine resources. *Mar Ecol Prog Ser* 274:269–303
- Chen YQ, Shen XQ (1999) Changes in the biomass of the East China Sea ecosystem. In: Sherman K, Tang Q (eds) Large Marine Ecosystems of the Pacific Rim: assessment, sustainability, and management. Blackwell Science, Malden, MA, p 221–239
- Duda AM (1982) Municipal point sources and agricultural nonpoint source contributions to coastal eutrophication. *Water Res Bull* 18(3):397–407
- Duda AM, El-Ashry MT (2000) Addressing the global water and environmental crises through integrated approaches to the management of land, water, and ecological resources. *Water Int* 25:115–126
- EPA (US Environmental Protection Agency) (2004) National coastal condition report II. Washington, DC, EPA-620/R-03/002
- FAO (1995) Code of conduct for responsible fisheries. FAO, Rome (available at: www.fao.org/Fl/agreem/codecond/ficonde.asp)
- Garcia SM, Zerbi A, Aliaume C, Do Chi T, Lassarre G (2003) The ecosystem approach to fisheries. Rep No. 443, FAO, Rome
- Garibaldi L, Limongelli L (2003). Trends in oceanic captures and clustering of large marine ecosystems: two studies based on the FAO capture database. *Fish Tech Pap* 435, FAO, Rome
- GESAMP (2001) The state of the marine environment. Group of Experts on the Scientific Aspects of Marine Pollution, Regional Seas Reports and Studies. UNEP, Nairobi
- Cislason H, Sinclair M (2000) Ecosystem effects of fishing. *ICES J Mar Sci* 57:465–791
- HELCOM (2001) Environment of the Baltic Sea area 1994–1998. *Baltic Sea Environment Proc* 82A, Helsinki Commission, Helsinki
- Howarth R, Anderson D, Cloern J, Elfring C and 7 others (2000) Nutrient pollution of coastal rivers, bays, and seas. *ESA Issues Ecol* 7:1–15
- Kroeze C, Seitzinger SP (1998) Nitrogen inputs to rivers, estuaries and continental shelves and related nitrous oxide emissions in 1990 and 2050: a global model. *Nutr Cycl Agroecosyst* 52:195–212
- Juda L, Hennessey T (2001) Governance profiles and the management of the uses of large marine ecosystems. *Ocean Dev Int Law* 32:41–67
- Levin SA (1990) Physical and biological scales and the modeling of predator-prey interactions in large marine ecosystems. In: Sherman K, Alexander LM, Gold BD (eds) Large marine ecosystems: patterns, processes and yields. American Association for the Advancement of Science, Washington, DC, p 179–187
- NMFS (National Marine Fisheries Service) (1999) Our living oceans. Report on the status of U.S. living marine resources, 1999. US Dep Commer, NOAA Tech Memo NMFS-F/SPO-41
- NRC (National Research Council) (2000) Clean coastal waters: understanding and reducing the effects of nutrient pollution. National Academy Press, Washington, DC
- Pauly D, Chuenpagdee R (2003) Development of fisheries in the Gulf of Thailand large marine ecosystem: analysis of an unplanned experiment. In: Hempel G, Sherman K (eds) Large Marine Ecosystems of the world: trends in exploitation, protection, and research. Elsevier, Amsterdam, p 337–354
- Perry AL, Low PJ, Ellis JR, Reynolds JD (2005) Climate change and distribution shifts in marine fishes. *Science* 308:1912–1915
- Pikitch EK, Santora C, Babcock EA, Bakun A and 13 others (2004) Ecosystem-based fishery management. *Science* 305:346–347
- Pitcher TJ (2001) Fisheries managed to rebuild ecosystems? Reconstructing the past to salvage the future. *Ecol Appl* 11:601–617
- Rabalais NN, Turner RE, Wiseman WJ Jr. (1999) Hypoxia in the northern Gulf of Mexico: linkages with the Mississippi River. In: Kumpf H, Steidinger K, Sherman K (eds) The Gulf of Mexico Large Marine Ecosystem: assessment, sustainability, and management. Blackwell Science, Malden, MA, p 297–322
- Rice J (2002) Changes to the Large Marine Ecosystem of the Newfoundland-Labrador Shelf. In: Sherman K, Skjoldal HR (eds) Large Marine Ecosystems of the North Atlantic: changing states and sustainability. Elsevier, Amsterdam, p 51–104
- Sainsbury K, Sumaila UR (2003) Incorporating ecosystem objectives into management of sustainable marine fisheries, including 'best practice' reference points and use of marine protected areas. In: Sinclair M, Valdimarsson G (eds) Responsible fisheries in the marine ecosystem. FAO, Rome & CABI Publishing, Wallingford, p 343–361
- Seitzinger SP, Kroeze C (1998) Global distribution of nitrous oxide production and N inputs to freshwater and coastal marine ecosystems. *Global Biogeochem Cycl* 12:93–113
- Sherman K (1994) Sustainability, biomass yields, and health of coastal ecosystems: an ecological perspective. *Mar Ecol Prog Ser* 112:277–301
- Sherman K, Kane J, Murawski S, Overholtz W, Solow A (2002) The US northeast shelf Large Marine Ecosystem: zooplankton trends in fish biomass recovery. In: Sherman K, Skjoldal HR (eds) Large Marine Ecosystems of the North Atlantic: changing states and sustainability. Elsevier, Amsterdam, p 195–215
- Sherman K, O'Reilly J, Kane J (2003) Assessment and sustainability of the U.S. northeast shelf ecosystem. In: Hempel G, Sherman K (eds) Large Marine Ecosystems of the world: trends in exploitation, protection, and research. Elsevier, Amsterdam, p 93–120
- Stergiou KI (2002) Overfishing, tropicalization of fish stocks, uncertainty and ecosystem management: resharpening Ockham's razor. *Fish Res* 55:109

-
- Sutinen JG (ed) (2000) A framework for monitoring and assessing socioeconomics and governance of large marine ecosystems. NOAA Tech Memo NMFS-NE-158
- Tang Q, Jin X (1999) Ecology and variability of economically important pelagic fishes in the Yellow Sea and Bohai Sea. In: Sherman K, Tang Q (eds) Large Marine Ecosystems of the Pacific Rim: assessment, sustainability, and management. Blackwell Science, Malden, MA, p 179–198
- Tang Q (2003) The Yellow Sea LME and mitigation action. In: Hempel G, Sherman K (eds) Large Marine Ecosystems of the world: trends in exploitation, protection, and research. Elsevier, Amsterdam, p 121–144
- Wang H (2004) An evaluation of the modular approach to the assessment and management of Large Marine Ecosystems. *Ocean Dev Int Law* 35:267–286
- Zwanenburg KCT (2003) The Scotian Shelf. In: Hempel G, Sherman K (eds) Large Marine Ecosystems of the world: trends in exploitation, protection, and research. Elsevier, Amsterdam, p 75–92
-